

In vivo release of high dose
vancomycin from loaded cement in
patients with periprostheses infection
effective bactericidal activity.

Delepine, N.; Alvarez, J.; Abe, E.;;
Markowska, B.;; Delepine, G.;



CENTRE HOSPITALIER
UNIVERSITAIRE
RAYMOND-POINCARÉ

104 Bd Raymond-Poincaré
92380 Garches
Standard : 01.47.10.79.00
contact-poincaré.garches
@rpc.ap-hop.paris.fr

After limb salvage, infection

- is a devastating complication that occurred in **8 to 20% of patients treated** by en bloc resection and prosthetic reconstruction for bone sarcomas.
- Resulting often in **secondary amputation**



Antibiotic loaded cement spacers

- Have been widely used since 1972 to prevent and to treat prosthetic infection
- The delivery of a high concentration of antibiotics in a localized area is thought to be safer than systemic administration of intravenous antibiotics in such doses

the emergence of increased resistance of staphylococcus



- Explains recent less effectiveness of conventional antibiotic loaded cement (low dose of antibiotics)
- Plaes for higher doses of antibiotics
- and compells us to consider the antibiotic concentration in the operating field

systemic safety of high dose vancomycin

- The systemic safety of high dose vancomycin loaded spacer has been investigated* but rarely the elution of high vancomycin from cement in vivo.

***Systemic Safety of High-Dose Antibiotic-Loaded Cement Spacers after Resection of an Infected Total Knee Arthroplasty** *Bryan D. Springer, MD and coll Clin. Orthop.Rel.Desease 2004*

The aims of the study

- 1^o) To **confirm the systemic safety** of using high doses of vancomycin in cement
- 2^o) To **evaluate the elution of vancomycin into the site** of the excision arthroplasty to see if effective bactericidal activity can be obtained

Patients and methods

- From 2006 to 2008 , 16 consecutive patients were managed by prosthetic exchange 2 stages procedure using high dose vancomycin loaded cement spacer.
- Patients were males :7 females : 9.
Average of age at the time of surgery was 22 years.

Antibiotic-loaded methylmethacrylate cement

- Cement were prepared by adding **4 g of vancomycin powder to a 40 g pack of Palacos R cement** in the operative room during the operation.




Antibiotic-loaded methylmethacrylate cement

- We generally used **2 to 4 batches of cement in one spacer** depending of the size and length of resection
- The **average dose of implanted vancomycin was 7.5 G (4-14.5)**.

spacer for proximal tibia

The spacer was composed of metallic rods covered with antibiotic loaded cement

Gentamycine+
Vancomycine (4 gr/pack).




spacers for femur



Distal femur



Total femur





Proximal femur



other spacers

Spacers used for proximal humerus or acetabular



Post operative cares

- The wounds were closed with absorbable mono-filaments sutures over one suction drain.
- **Intravenous antibiotics excluding vancomycin were given for 6 to 24 weeks.**
- **Patients biological values and the concentrations of vancomycin in the blood and in the aliquots of suction drainage were checked daily until removal of drain (d10-d15).**



Results for systemic safety


- **The serum concentration of vancomycin remained under 2 $\mu\text{g/ml}$ in all patients**
- We observed no case of
 - allergy,
 - toxicity
 - or intolerance

Local concentration

- Local concentration of vancomycin depended **of the dose of vancomycin used** and decreased quickly during the first week
- half life :2.25 days.



Local concentration for 10 Grams



- For a dose of 10 G vancomycin , the average concentration from the drain was :d1 :725 $\mu\text{g/ml}$, d2 :510 $\mu\text{g/ ml}$,d3 :346

Is it bactericidal?

- These results should be compared to the bactericidal concentration of vancomycin **for staphylococcus aureus** :
- **10 to 20 $\mu\text{g/ml}$ for usual organisms,**
- **20 to 40 $\mu\text{g/ml}$ for resistant organisms .**



Conclusion

- high dose vancomycin spacers result in very low serum concentration without risk of systemic toxicity.
- In the operative wound , very high concentration are obtained , 10 to 20 fold bactericidal concentration for staphylococcus aureus.
- Additional studies are needed , with longer follow-up to evaluate the clinical efficacy of this method.